arXiv:1703.07664v1 [math.GM] 7 Mar 2017

ASSESMENT OF SMARANDACHE CURVES IN THE NULL CONE $\mathbf{Q}^2$

MIHRIBAN KULAHCI AND FATMA ALMAZ

Abstract. In this paper, we give Smarandache curves according to the asymptotic orthonormal frame in null cone $\mathbf{Q}^2$. By using cone frame formulas, we present some characterizations of Smarandache curves and calculate cone frenet invariants of these curves. Also, we illustrate these curves with an example.

1. Introduction

Human being were bewitched by curves and curved shapes long before they took into account them as mathematical objects. But the greatest effect in the research of curves was, of course, the discovery of the calculus. Geometry before calculus includes only the simplest curves.

In classical curve theory, the geometry of a curve in three-dimensions is actually characterized by Frenet vectors.

Smarandache geometry is a geometry which has at least one Smarandachely denied axiom [4]. An axiom is said to be Smarandachely denied, if it behaves in at least two different ways within the same space. Smarandache curve is defined as a regular curve whose position vector is composed by Frenet frame vectors of another regular curve. Smarandache curves in various ambient spaces have been classfied in [1]-[14], [19], [21]-[32].

In this study, we define special Smarandache curves such as $x\alpha, xy, \alpha y$ and $x\alpha y$ Smarandache curves according to asymptotic orthonormal frame in the null cone $\mathbf{Q}^2$ and we examine the curvature and the asymptotic orthonormal frame's vectors of Smarandache curves. We also give an example related to these curves.

2. Preliminaries

Some basics of the curves in the null cone are provided from, [15]-[18].

Let E_1^3 be the 3−dimensional pseudo-Euclidean space with the

$$\widetilde{g}(X,Y) = \langle X, Y\rangle = x_1y_1 + x_2y_2 - x_3y_3$$

for all $X = (x_1, x_2, x_3)$, $Y = (y_1, y_2, y_3) \in E_1^3$. E_1^3 is a flat pseudo-Riemannian manifold of signature $(2, 1)$.

Date: 2017.

2000 *Mathematics Subject Classification.* Primary 53A40; Secondary 53A35.

Key words and phrases. Smarandache curve, asymptotic orthonormal frame, null cone, cone frame formulas.

This paper is in final form and no version of it will be submitted for publication elsewhere.

Let M be a submanifold of E_1^3. If the pseudo-Riemannian metric $\widetilde{g}$ of E_1^3 induces a pseudo-Riemannian metric g(respectively, a Riemannian metric, a degenerate quadratic form) on M, then M is called a timelike(respectively, spacelike, degenerate) submanifold of E_1^3. Let c be a fixed point in E_1^3. The pseudo-Riemannian lightlike cone(quadric cone) is defined by

$$\mathbf{Q}_1^2(c) = \left\{x \in E_1^3 : g(x-c, x-c) = 0\right\},$$

where the point c is called the center of $\mathbf{Q}_1^2(c)$. When $c = 0$, we simply denote $\mathbf{Q}_1^2(0)$ by $\mathbf{Q}^2$ be and call it the null cone.

Let E_1^3 be 3-dimensional Minkowski space and $\mathbf{Q}^2$ the lightlike cone in E_1^3. A vector $V \neq 0$ in E_1^3 is called spacelike, timelike or lightlike, if $\langle V, V\rangle > 0$, $\langle V, V\rangle < 0$ or $\langle V, V\rangle = 0$, respectively. The norm of a vector $x \in E_1^3$ is given by $\|x\| = \sqrt{\langle x, x\rangle}$, [20].

We assume that curve $x : I \to \mathbf{Q}^2 \subset E_1^3$ is a regular curve in $\mathbf{Q}^2$ for $t \in I$. In the following, we always assume that the curve is regular.

A frame field $\{x, \alpha, y\}$ on E_1^3 is called an asymptotic orthonormal frame field, if

$$\langle x, x\rangle = \langle y, y\rangle = \langle x, \alpha\rangle = \langle y, \alpha\rangle = 0,\ \langle x, y\rangle = \langle \alpha, \alpha\rangle = 1.$$

Using $x'(s) = \alpha(s)$ we know that $\{x(s), \alpha(s), y(s)\}$ from an asymptotic orthonormal frame along the curve $x(s)$ and the cone frenet formulas of $x(s)$ are given by

$$\begin{aligned} x'(s) &= \alpha(s) \\ \alpha'(s) &= \kappa(s)x(s) - y(s) \\ y'(s) &= -\kappa(s)\alpha(s) \end{aligned} \tag{2.1}$$

where the function $\kappa(s)$ is called cone curvature function of the curve $x(s)$, [17].

Let $x : I \to \mathbf{Q}^2 \subset E_1^3$ be a spacelike curve in $\mathbf{Q}^2$ with an arc length parameter s. Then $x = x(s) = (x_1, x_2, x_3)$ can be written as

$$x(s) = \frac{f_s^{-1}}{2}(f^2 - 1, 2f, f^2 + 1) \tag{2.2}$$

for some non constant function $f(s)$ and $f_s = f'$, [18].

3. Smarandache Curves in The Null Cone $\mathbf{Q}^2$

In this section, we define the Smarandache curves according to the asymptotic orthonormal frame in $\mathbf{Q}^2$. Also, we obtain the asymptotic orthonormal frame and cone curvature function of the Smarandache partners lying on $\mathbf{Q}^2$ using cone frenet formulas.

Smarandache curve $\gamma = \gamma(s^*(s))$ of the curve x is a regular unit speed curve lying fully on $\mathbf{Q}^2$. Let $\{x, \alpha, y\}$ and $\{\gamma, \alpha_\gamma, y_\gamma\}$ be the moving asymptotic orthonormal frames of x and γ, respectively.

Definition 1. *Let x be unit speed spacelike curve lying on $\mathbf{Q}^2$ with the moving asymptotic orthonormal frame $\{x, \alpha, y\}$. Then, $x\alpha-$smarandache curve of x is defined by*

$$\gamma_{x\alpha}(s^*) = \frac{c}{b}x(s) + \alpha(s), \tag{3.1}$$

where $c, b \in \mathbb{R}_0^+$.

Theorem 1. *Let x be unit speed spacelike curve in $\mathbf{Q}^2$ with the moving asymptotic orthonormal frame $\{x, \alpha, y\}$ and cone curvature $\kappa(s)$ and let $\gamma_{x\alpha}$ be $x\alpha-$smarandache curve with asymptotic orthonormal frame $\{\gamma_{x\alpha}, \alpha_{x\alpha}, y_{x\alpha}\}$. Then the following relations hold:*

i) The asymptotic orthonormal frame $\{\gamma_{x\alpha}, \alpha_{x\alpha}, y_{x\alpha}\}$ of the $x\alpha$-smarandache curve $\gamma_{x\alpha}$ is given as

$$\begin{bmatrix} \gamma_{x\alpha} \\ \alpha_{x\alpha} \\ y_{x\alpha} \end{bmatrix} = \begin{bmatrix} \frac{c}{b} & 1 & 0 \\ \psi & \frac{b}{c}\kappa\psi & -\frac{b}{c}\psi \\ \varrho_1 & \varrho_2 & \varrho_3 \end{bmatrix} \begin{bmatrix} x \\ \alpha \\ y \end{bmatrix} \tag{3.2}$$

where

$$\psi = \frac{c}{\sqrt{c^2 - 2\kappa b^2}}, \tag{3.3}$$

$$\Upsilon_1 = \frac{b}{c}(\frac{b}{c}\kappa' + \kappa)\psi\psi' + \frac{b^2}{c^2}\kappa\left(\psi'\right)^2,$$

$$\Upsilon_2 = \frac{b}{c}\psi(\psi' + 2\frac{b}{c}\kappa\psi), \tag{3.4}$$

$$\Upsilon_3 = -\frac{b}{c}\psi(\frac{b}{c}\psi' + \psi)$$

and

$$\begin{aligned} \varrho_1 &= -(\Upsilon_1 + \frac{c}{2b}\left(2\Upsilon_1\Upsilon_3 + \Upsilon_2^2\right)) = -\Upsilon_1 + \frac{c}{b}\kappa_{\gamma_{x\alpha}}(s^*), \\ \varrho_2 &= -(\Upsilon_2 + \frac{1}{2}\left(2\Upsilon_1\Upsilon_3 + \Upsilon_2^2\right)) = -\Upsilon_2 + \kappa_{\gamma_{x\alpha}}(s^*), \\ \varrho_3 &= -\Upsilon_3. \end{aligned} \tag{3.5}$$

ii) The cone curvature $\kappa_{\gamma_{x\alpha}}(s^*)$ of the curve $\gamma_{x\alpha}$ is given by

$$\kappa_{\gamma_{x\alpha}}(s^*) = -\frac{1}{2}\left(2\Upsilon_1\Upsilon_3 + \Upsilon_2^2\right). \tag{3.6}$$

where

$$s^* = \frac{1}{b}\int \sqrt{c^2 - 2b^2\kappa(s)}ds.$$

Proof. **i)** We assume that the curve x is a unit speed spacelike curve with the asymptotic orthonormal frame $\{x, \alpha, y\}$ and cone curvature κ. Differentiating the equation (3.1) with respect to s and considering (2.1), we have

$$\gamma'_{x\alpha}(s^*) = A\left(\alpha(s) + \frac{b}{c}\kappa x(s) - \frac{b}{c}y(s)\right), \tag{3.7}$$

where

$$\frac{ds^*}{ds} = \frac{1}{b}\sqrt{c^2 - 2b^2\kappa(s)}, \tag{3.8}$$

$$\psi = \frac{c}{\sqrt{c^2 - 2\kappa(s)b^2}}. \tag{3.9}$$

It can be easily seen that the tangent vector $\gamma'_{x\alpha}(s^*) = \alpha_{x\alpha}(s^*)$ is a unit spacelike vector.

Differentiating (3.7) , we obtain equation as follows

$$\gamma''_{x\alpha}(s^*) = \Upsilon_1 x(s) + \Upsilon_2\alpha(s) + \Upsilon_3 y(s), \tag{3.10}$$

where

$$\Upsilon_1 = \frac{b}{c}(\frac{b}{c}\kappa' + \kappa)\psi\psi' + \frac{b^2}{c^2}\kappa\left(\psi'\right)^2,$$
$$\Upsilon_2 = \frac{b}{c}\psi(\psi' + 2\frac{b}{c}\kappa\psi),$$
$$\Upsilon_3 = -\frac{b}{c}\psi(\frac{b}{c}\psi' + \psi).$$

$$y_{x\alpha}(s^*) = -\gamma''_{x\alpha} - \frac{1}{2}\left\langle \gamma''_{x\alpha}, \gamma''_{x\alpha}\right\rangle \gamma_{x\alpha} \tag{3.11}$$

By the help of previous equation (3.11), we obtain

$$y_{x\alpha}(s^*) = \varrho_1 x(s) + \varrho_2\alpha(s) + \varrho_3 y(s), \tag{3.12}$$

where $\varrho_1 = -(\Upsilon_1 + \frac{c}{2b}\left(2\Upsilon_1\Upsilon_3 + \Upsilon_2^2\right))$, $\varrho_2 = -(\Upsilon_2 + \frac{1}{2}\left(2\Upsilon_1\Upsilon_3 + \Upsilon_2^2\right))$, $\varrho_3 = -\Upsilon_3$.

ii) The curvature $\kappa_{\gamma_{x\alpha}}(s^*)$ of the $\gamma_{x\alpha}(s^*)$ is explicity obtained by

$$\begin{aligned}\kappa_{\gamma_{x\alpha}}(s^*) &= -\frac{1}{2}\left\langle \gamma''_{x\alpha}, \gamma''_{x\alpha}\right\rangle \\ &= -\frac{1}{2}\left(2\Upsilon_1\Upsilon_3 + \Upsilon_2^2\right).\end{aligned} \tag{3.13}$$

Thus, the theorem is proved. □

Definition 2. *Let x be unit speed spacelike curve lying on $\mathbf{Q}^2$ with the moving asymptotic orthonormal frame $\{x, \alpha, y\}$. Then, $xy-$smarandache curve of x is defined by*

$$\gamma_{xy}(s^*) = \frac{1}{\sqrt{2cb}}\left(cx(s) + by(s)\right), \tag{3.14}$$

where $c, b \in \mathbb{R}_0^+$.

Theorem 2. *Let x be unit speed spacelike curve in $\mathbf{Q}^2$ with the moving asymptotic orthonormal frame $\{x, \alpha, y\}$ and cone curvature κ and let γ_{xy} be $xy-$smarandache curve with asymptotic orthonormal frame $\{\gamma_{xy},\ \alpha_{xy},\ y_{xy}\}$. Then the following relations hold:*

i) The asymptotic orthonormal frame $\{\gamma_{xy}, \alpha_{xy}, y_{xy}\}$ of the $xy-$smarandache curve γ_{xy} is given as

$$\begin{bmatrix}\gamma_{xy}\\ \alpha_{xy}\\ y_{xy}\end{bmatrix} = \begin{bmatrix}\sqrt{\frac{c}{2b}} & 0 & \sqrt{\frac{b}{2c}}\\ 0 & 1 & 0\\ \frac{b\kappa^2\sqrt{2bc}}{(c-b\kappa)^2} & 0 & \frac{c\sqrt{2bc}}{(c-b\kappa)^2}\end{bmatrix}\begin{bmatrix}x\\ \alpha\\ y\end{bmatrix}. \tag{3.15}$$

ii) The cone curvature $\kappa_{\gamma_{xy}}(s^*)$ of the curve γ_{xy} is given by

$$\kappa_{\gamma_{xy}}(s^*) = \frac{2bc\kappa(s)}{\left(c - b\kappa(s)\right)^2}, \tag{3.16}$$

where

$$s^* = \frac{1}{\sqrt{2cb}}\int \left(c - b\kappa(s)\right) ds. \tag{3.17}$$

Proof. **i)** We assume that the curve x is a unit speed spacelike curve with the asymptotic orthonormal frame $\{x, \alpha, y\}$ and cone curvature κ. Differentiating the equation (3.14) with respect to s and considering (2.1), we have

$$\gamma'_{xy}(s^*)\frac{ds^*}{ds} = \frac{1}{\sqrt{2cb}}\left(c - b\kappa(s)\right)\overrightarrow{\alpha}(s). \tag{3.18}$$

By considering (3.17), we get

$$\gamma'_{xy}(s^*) = \alpha(s) = \alpha_{xy}. \tag{3.19}$$

Here, it can be easily seen that the tangent vector $\overrightarrow{\alpha}_{xy}$ is a unit spacelike vector.

$$\gamma''_{xy}(s^*)\frac{ds^*}{ds} = \kappa x(s) - y(s). \tag{3.20}$$

By substituting (3.17) into (3.20) and making necessary calculations, we obtain

$$\gamma''_{xy}(s^*) = \frac{\kappa\sqrt{2bc}}{(c - b\kappa)}\overrightarrow{x} - \frac{\sqrt{2bc}}{(c - b\kappa)^2}\overrightarrow{y}. \tag{3.21}$$

By the help of equation $y_{xy}(s^*) = -\gamma''_{xy} - \frac{1}{2}\left\langle \gamma''_{xy}, \gamma''_{xy}\right\rangle \gamma_{xy}$, we write

$$y_{xy}(s^*) = \frac{b\sqrt{2bc}.\kappa^2}{(c - b\kappa)^2}x(s) + \frac{c\sqrt{2bc}}{(c - b\kappa)^2}y(s). \tag{3.22}$$

ii) The curvature $\kappa_{\gamma_{xy}}(s^*)$ of the $\gamma_{xy}(s^*)$ is explicity obtained by

$$\kappa_{\gamma_{xy}}(s^*) = -\frac{1}{2}\left\langle \gamma''_{xy}, \gamma''_{xy}\right\rangle = \frac{2bc\kappa(s)}{(c - b\kappa(s))^2}.$$

□

Definition 3. *Let x be unit speed spacelike curve lying on $\mathbf{Q}^2$ with the moving asymptotic orthonormal frame $\{x, \alpha, y\}$. Then, $\alpha y-$smarandache curve of x is defined by*

$$\gamma_{\alpha y}(s^*) = \alpha(s) + \frac{b}{c}y(s), \tag{3.23}$$

where $c, b \in \mathbb{R}_0^+$.

Theorem 3. *Let x be unit speed spacelike curve in $\mathbf{Q}^2$ with the moving asymptotic orthonormal frame $\{x, \alpha, y\}$ and cone curvature κ and let $\gamma_{\alpha y}$ be $\alpha y-$smarandache curve with asymptotic orthonormal frame $\{\gamma_{\alpha y},\ \alpha_{\alpha y},\ y_{\alpha y}\}$. Then the following relations hold:*

i) The asymptotic orthonormal frame $\{\gamma_{\alpha y}, \alpha_{\alpha y}, y_{\alpha y}\}$ of the $\alpha y-$smarandache curve $\gamma_{\alpha y}$ is given as

$$\begin{bmatrix}\gamma_{\alpha y}\\ \alpha_{\alpha y}\\ y_{\alpha y}\end{bmatrix} = \begin{bmatrix} 0 & 1 & \frac{b}{c}\\ \frac{c\sqrt{\kappa}}{\sqrt{b^2-2c^2}} & \frac{b\sqrt{\kappa}}{\sqrt{b^2-2c^2}} & \frac{c\sqrt{\kappa}}{\kappa\sqrt{b^2-2c^2}}\\ \omega_1 & \omega_2 & \omega_3\end{bmatrix}\begin{bmatrix}x\\ \alpha\\ y\end{bmatrix}, \tag{3.24}$$

where

$$\begin{aligned}\zeta_1 &= \frac{c\kappa'}{b^2-2c^2}\left(\frac{c-2\kappa b}{2\kappa}\right),\\ \zeta_2 &= \frac{c\kappa'}{b^2-2c^2}\left(\frac{c-b-1}{2\kappa}\right),\\ \zeta_3 &= \frac{c\kappa'}{b^2-2c^2}\left(\frac{b\kappa-c}{2\kappa^2}\right)\end{aligned} \tag{3.25}$$

and

$$\begin{aligned}\omega_1 &= -\zeta_1,\\ \omega_2 &= -(\zeta_2+\frac{1}{2}\left(2\zeta_1\zeta_3+\zeta_2^2\right)),\\ \omega_3 &= -(\zeta_3+\frac{b}{2c}\left(2\zeta_1\zeta_3+\zeta_2^2\right)).\end{aligned} \tag{3.26}$$

ii) The cone curvature $\kappa_{\gamma_{\alpha y}}(s^*)$ of the curve $\gamma_{\alpha y}$ is given by

$$\kappa_{\gamma_{\alpha y}}(s^*) = -\frac{c^2}{8\left(b^2-2c^2\right)^2}(\frac{\kappa'}{\kappa})^2\left(\frac{1}{\kappa}\left(c-2\kappa b\right)\left(b\kappa-c\right)+\left(c-b-1\right)^2\right), \tag{3.27}$$

where

$$s^* = \frac{\sqrt{b^2-2c^2}}{c}\int\sqrt{\kappa(s)}ds. \tag{3.28}$$

Proof. **i)** Let the curve x be a unit speed spacelike curve with the asymptotic orthonormal frame $\{x,\alpha,y\}$ and cone curvature κ. Differentiating the equation (3.23) with respect to s and considering (2.1), we find

$$\gamma'_{\alpha y}(s^*)\frac{ds^*}{ds} = \kappa\overrightarrow{x(s)} - \frac{b}{c}\kappa\overrightarrow{\alpha(s)} - \overrightarrow{y(s)}.$$

This can be written as following

$$\alpha_{\alpha y}(s^*)\frac{ds^*}{ds} = \kappa\overrightarrow{x(s)} - \frac{b}{c}\kappa\overrightarrow{\alpha(s)} - \overrightarrow{y(s)}, \tag{3.29}$$

where

$$\frac{ds^*}{ds} = \frac{\sqrt{b^2-2c^2}}{c}\sqrt{\kappa(s)}. \tag{3.30}$$

By substituting (3.30) into (3.29), we find

$$\alpha_{\alpha y}(s^*) = \frac{c\sqrt{\kappa}}{\sqrt{b^2-2c^2}}\overrightarrow{x} - \frac{b\sqrt{\kappa}}{\sqrt{b^2-2c^2}}\overrightarrow{\alpha} - \frac{c\sqrt{\kappa}}{\kappa\sqrt{b^2-2c^2}}\overrightarrow{y}. \tag{3.31}$$

Differentiating (3.31) and using (3.30), we get

$$\gamma''_{\alpha y}(s^*) = \zeta_1 x(s) + \zeta_2\alpha(s) + \zeta_3 y(s),$$

where $\zeta_1 = \frac{c\kappa'}{b^2-2c^2}\left(\frac{c-2\kappa b}{2\kappa}\right), \zeta_2 = \frac{c\kappa'}{b^2-2c^2}\left(\frac{c-b-1}{2\kappa}\right), \zeta_3 = \frac{c\kappa'}{b^2-2c^2}\left(\frac{b\kappa-c}{2\kappa^2}\right).$

$$y_{\alpha y}(s^*) = -\gamma''_{\alpha y} - \frac{1}{2}\left\langle\gamma''_{\alpha y},\gamma''_{\alpha y}\right\rangle\gamma_{\alpha y}. \tag{3.32}$$

By the help of equation (3.32), we obtain

$$y_{\alpha y}(s^*) = \omega_1 x(s) + \omega_2\alpha(s) + \omega_3 y(s), \tag{3.33}$$

where $\omega_1 = -\zeta_1$, $\omega_2 = -(\zeta_2+\frac{1}{2}\left(2\zeta_1\zeta_3+\zeta_2^2\right))$, $\omega_3 = -(\zeta_3+\frac{b}{2c}\left(2\zeta_1\zeta_3+\zeta_2^2\right))$.

ii) The curvature $\kappa_{\gamma_{\alpha y}}(s^*)$ of the $\gamma_{\alpha y}(s^*)$ is explicity obtained by

$$\kappa_{\gamma_{\alpha y}}(s^*) = -\frac{c^2}{8\left(b^2-2c^2\right)^2}\left(\frac{\kappa'}{\kappa}\right)^2\left(\frac{(c-2\kappa b)(b\kappa-c)}{\kappa}+(c-b-1)^2\right).$$

□

Definition 4. *Let x be unit speed spacelike curve lying on $\mathbf{Q}^2$ with the moving asymptotic orthonormal frame $\{x,\alpha,y\}$. Then, $x\alpha y-$smarandache curve of x is defined by*

$$\gamma_{x\alpha y}(s^*) = \frac{1}{\sqrt{2cc^*+b^2}}\left(cx(s)+b\alpha(s)+c^*y(s)\right), \tag{3.34}$$

where $c, c^*, b \in \mathbb{R}_0^+$.

Theorem 4. *Let x be unit speed spacelike curve in $\mathbf{Q}^2$ with the moving asymptotic orthonormal frame $\{x,\alpha,y\}$ and cone curvature κ and let $\gamma_{x\alpha y}$ be $x\alpha y-$smarandache curve with asymptotic orthonormal frame $\{\gamma_{x\alpha y},\ \alpha_{x\alpha y},\ y_{x\alpha y}\}$. Then the following relations hold:*

i) The asymptotic orthonormal frame $\{\gamma_{x\alpha y},\ \alpha_{x\alpha y},\ y_{x\alpha y}\}$ of the $x\alpha y-$smarandache curve $\gamma_{x\alpha y}$ is given as

$$\begin{bmatrix}\gamma_{x\alpha y}\\ \alpha_{x\alpha y}\\ y_{x\alpha y}\end{bmatrix} = \begin{bmatrix}\frac{c}{\sqrt{2cc^*+b^2}} & \frac{b}{\sqrt{2cc^*+b^2}} & \frac{c^*}{\sqrt{2cc^*+b^2}}\\ \rho_1 & \rho_2 & \rho_3\\ \sigma_1 & \sigma_2 & \sigma_3\end{bmatrix}\begin{bmatrix}x\\ \alpha\\ y\end{bmatrix} \tag{3.35}$$

where

$$\begin{aligned}&\eta = \sqrt{\left(c-c^*\kappa(s)\right)^2-2b^2\kappa(s)}\\ &\rho_1 = \frac{b\kappa(s)}{\eta}, \rho_2 = \frac{c-c^*\kappa(s)}{\eta}, \rho_3 = -\frac{b}{\eta}\\ &\xi_1 = \left(\rho_1'+\rho_2\kappa\right), \xi_2 = \rho_2'+\rho_1+\rho_3\kappa, \xi_3 = -\rho_3'-\rho_2\end{aligned} \tag{3.36}$$

and

$$\begin{aligned}\sigma_1 &= -\xi_1 - \frac{c}{2\sqrt{2cc^*+b^2}}\left(2\xi_1\xi_3+\xi_2^2\right),\\ \sigma_2 &= -\xi_2 - \frac{b}{2\sqrt{2cc^*+b^2}}\left(2\xi_1\xi_3+\xi_2^2\right),\\ \sigma_3 &= -\xi_3 - \frac{c^*}{2\sqrt{2cc^*+b^2}}\left(2\xi_1\xi_3+\xi_2^2\right).\end{aligned} \tag{3.37}$$

ii) The cone curvature $\kappa_{\gamma_{x\alpha y}}(s^*)$ of the curve $\gamma_{x\alpha y}$ is given by

$$\begin{aligned}\kappa_{\gamma_{yx\alpha}}(s^*) = &\left(b\left(\frac{\kappa}{\eta}\right)'+\frac{c-c^*\kappa}{\eta}\kappa\right)\left(\left(\frac{b}{\eta}\right)'+\frac{c-c^*\kappa}{\eta}\right)\\ &-\frac{1}{2}\left(\frac{c-c^*\kappa}{\eta}\right)'.\end{aligned} \tag{3.38}$$

where

$$s^* = \frac{1}{\sqrt{2cc^*+b^2}}\int\sqrt{\left(c-c^*\kappa(s)\right)^2-2b^2\kappa(s)}ds,\ b,c,c^*\in\mathbb{R}_0^+. \tag{3.39}$$

Proof. **i)** Differentiating the equation (3.34) with respect to s and considering (2.1), we find

$$\gamma'_{x\alpha y}(s^*)\frac{ds^*}{ds} = \frac{1}{\sqrt{2cc^*+b^2}}\left(b\kappa\overrightarrow{x(s)} + (c-c^*\kappa)\overrightarrow{\alpha(s)} - \overrightarrow{by(s)}\right). \tag{3.40}$$

This can be written as follows

$$\alpha_{x\alpha y}(s^*) = \frac{b\kappa}{\eta}\overrightarrow{x(s)} + \frac{c-c^*\kappa}{\eta}\overrightarrow{\alpha(s)} - \frac{b}{\eta}\overrightarrow{y(s)}, \tag{3.41}$$

or

$$\alpha_{x\alpha y}(s^*) = \rho_1\overrightarrow{x(s)} + \rho_2\overrightarrow{\alpha(s)} - \rho_3\overrightarrow{y(s)}. \tag{3.42}$$

where

$$\frac{ds^*}{ds} = \frac{1}{\sqrt{2cc^*+b^2}}\sqrt{(c-c^*\kappa)^2 - 2b^2\kappa} \tag{3.43}$$

Differentiating (3.42) and using (3.43), we get

$$\gamma''_{xy\alpha}(s^*) = \xi_1 x(s) + \xi_2\alpha(s) + \xi_3 y(s),$$

where $\xi_1 = (\rho_1' + \rho_2\kappa)$, $\xi_2 = \rho_2' + \rho_1 + \rho_3\kappa$, $\xi_3 = -\rho_3' - \rho_2$.

$$y_{x\alpha y}(s^*) = -\gamma''_{x\alpha y} - \frac{1}{2}\left\langle\gamma''_{x\alpha y}, \gamma''_{x\alpha y}\right\rangle\gamma_{x\alpha y}. \tag{3.44}$$

By the help of equation (3.44), we obtain

$$y_{xay}(s^*) = \sigma_1 x(s) + \sigma_2\alpha(s) + \sigma_3 y(s), \tag{3.45}$$

where $\sigma_1 = -\xi_1 - \frac{c}{2\sqrt{2cc^*+b^2}}\left(2\xi_1\xi_3 + \xi_2^2\right)$, $\sigma_2 = -\xi_2 - \frac{b}{2\sqrt{2cc^*+b^2}}\left(2\xi_1\xi_3 + \xi_2^2\right)$, $\sigma_3 = -\xi_3 - \frac{c^*}{2\sqrt{2cc^*+b^2}}\left(2\xi_1\xi_3 + \xi_2^2\right)$.

ii) From $\kappa_{\gamma_{x\alpha y}}(s^*) = -\frac{1}{2}\left\langle\gamma''_{x\alpha y}, \gamma''_{x\alpha y}\right\rangle$, we have

$$\kappa_{\gamma_{yx\alpha}}(s^*) = \left(b\left(\frac{\kappa}{\eta}\right)' + \frac{c-c^*\kappa}{\eta}\kappa\right)\left(\left(\frac{b}{\eta}\right)' + \frac{c-c^*\kappa}{\eta}\right) - \frac{1}{2}\left(\frac{c-c^*\kappa}{\eta}\right)'.$$

□

Theorem 5. *Let $x : I \to \mathbf{Q}^2 \subset E_1^3$ be a spacelike curve in $\mathbf{Q}^2$ as follows*

$$x(s) = \frac{f_s^{-1}}{2}(f^2-1, 2f, f^2+1), \tag{3.46}$$

for some non constant function $f(s)$. Then we can write the following statements:

- *If x is a $x\alpha-$ smarandache curve, then the $x\alpha-$ smarandache curve $\gamma_{x\alpha}$ can be written as*

$$\gamma_{x\alpha}(s^*) = (\frac{c}{b} - f_s^{-1}f_{ss})x(s) + (f, 1, f). \tag{3.47}$$

- *If x is a $xy-$ smarandache curve, then the $xy-$ smarandache curve γ_{xy} can be written as*

$$\gamma_{xy}(s^*) = \frac{1}{\sqrt{2bc}}\left((c - \frac{1}{2}f_s^{-2}f_{ss}^2)x(s) + f_s^{-1}f_{ss}(f, 1, f) - f_s(1.0, 1)\right). \tag{3.48}$$

- *If x is a $\alpha y-$ smarandache curve, then the $\alpha y-$ smarandache curve $\gamma_{\alpha y}$ can be written as*

$$\gamma_{\alpha y}(s^*) = \left(-f_s^{-1}f_{ss} - \frac{b}{2c}f_s^{-2}f_{ss}^2\right)x(s) + \left(1 + \frac{b}{c}f_s^{-1}f_{ss}\right)(f,1,f) - \frac{b}{c}f_s\,(1.0,1)\,. \tag{3.49}$$

- *If x is a $x\alpha y-$ smarandache curve, then the $x\alpha y-$ smarandache curve $\gamma_{x\alpha y}$ can be written as*

$$\gamma_{x\alpha y}(s^*) = \frac{1}{\sqrt{2cc^* + b^2}}\begin{pmatrix} \left(c - bf_s^{-1}f_{ss} - \frac{c^*}{2}f_s^{-2}f_{ss}^2\right)x(s) \\ +\left(b + c^*f_s^{-1}f_{ss}\right)(f,1,f) - f_s\,(1.0,1) \end{pmatrix},$$

where $c, b, c^* \in \mathbb{R}_0^+$.

Proof. It is obvious from (3.1), (3.14), (3.23), (3.34) and (3.46). □

We can give the following example to hold special Smarandache curves in the null cone $\mathbf{Q}^2$. $x\alpha$, xy, αy, and $x\alpha y$ special smarandache curves of x curves are given in Figure 1 A, C, E, G, I, respectively. These figures rotated in three dimensions are also given in Figure 1 B, D, F, H, J, respectively.

Example 1. *The curve*

$$x(s) = \left(\frac{\cosh s}{2} - \frac{1}{\cosh s}, \tanh s, \frac{\cosh s}{2}\right)$$

is spacelike in $\mathbf{Q}^2$ with arc length parameter s. Also, the shape of the x curve is given as follows

Then we can write the smarandache curves of the x-curve as follows:

i) $x\alpha-$ smarandache curve $\gamma_{x\alpha}$ is given by

$$\gamma_{x\alpha}(s) = \begin{pmatrix} d\left(\frac{\cosh s}{2} - \frac{1}{\cosh s}\right) + \frac{\sinh s}{2} - \frac{\tanh s}{\cosh s}, \\ d\tanh s + \frac{1}{\cosh^2 s}, \\ d\frac{\cosh s}{2} + \frac{\sinh s}{2} \end{pmatrix}$$

ii) $xy-$ smarandache curve γ_{xy} is given by

$$\gamma_{xy}(s) = \begin{pmatrix} m\cosh s - n\tanh s\sinh s - d\left(\frac{1c + \tanh^2 s}{\cosh s}\right), \\ c\tanh s\left(e - \frac{\tanh^2 s}{2}\right), \\ d\left(\left(c - \frac{\tanh^2 s}{2}\right)\frac{\cosh s}{2} - \frac{1}{\cosh s}\right) \end{pmatrix}$$

iii) $\alpha y-$ smarandache curve $\gamma_{\alpha y}$ is given by

$$\gamma_{\alpha y}(s) = \begin{pmatrix} \left(\frac{\cosh s}{2} - \frac{1}{\cosh s}\right)(1 - e\tanh s)\tanh s + \sinh s\,(1 + d\tanh s) \\ -d\cosh s, \\ -\tanh^2 s\,(1 + d\tanh s) + d\tanh s, \\ \frac{\sinh s}{2} - d\cosh s - e\sinh s\tanh s \end{pmatrix}$$

iv) $x\alpha y-$ smarandache curve $\gamma_{x\alpha y}$ is given by

$$\gamma_{x\alpha y}(s) = \begin{pmatrix} m\sinh s - n\cosh s + d\sinh s\tanh s - \frac{c}{\cosh s} - \frac{b\sinh s - c^*\tanh^2 s}{\cosh^2 s}, \\ m + n\tanh s + k\tanh^2 s + l\tanh^3 s, \\ m\cosh s + d\sinh s + c\tanh s\sinh s, \end{pmatrix}$$

where $m, n, d, c, k, l, e \in \mathbb{R}_0^+$.



(A) The curve x



(B) The rotated surface of curve x



(C) $x\alpha-$smarandache curve



(D) $x\alpha$- smarandache surface.



(E) $xy-$smarandache curve



(F) $xy-$smarandache surface.



(G) αy$-$smarandache curve.



(H) αy$-$smarandache surface.



(I) $x\alpha$y$-$smarandache curve.



(J) $x\alpha$y$-$smarandache surface.

FIGURE 1

References

[1] Abdel-Aziz, H.S., Saad, M.K., Smarandache Curves of Some Special Curves in the Galilean 3−Space, Honan Mathematical J., 37(2), 253-264, 2015.

[2] Ali, A.T., " Special Smarandache Curves in the Euclidean Space", International Journal of Mathematical Combinatorics, vol.2, 30-36, 2010.

[3] Ali, A.T., Time-like Smarandache Curves Derived from a Space-like Helix, Journal of Dynamical Systems and Geometric Theories, 8(1), 93-100, 2010.

[4] Ashbacher, C., Smarandache Geometries, Smarandache Notions Journal, 8(1-3), 212-215, 1997.

[5] Bayrak, N., Bektas, O. and Yuce, S., Special Smarandache Curves in $\mathbb{R}_1^3$, Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat., 65(2), 143-160, 2016.

[6] Bektas, O. and Yuce, S., "Special Smarandache Curves According to Darboux Frame in E^3", Romanian Journal of Mathematics and Computer Science, 3(1),48-59, 2013.

[7] Cetin, M., Kocayiğit, H., On the Quaternionic Smarandache Curves in Euclidean 3−Space, 8(3),139-150, 2013.

[8] Cetin, M., Tuncer, Y., Karacan, M.K., Smarandache Curves According to Bishop Frame in Euclidean 3−Space, Gen. Math. Notes, 20(2), 50-66, 2014.

[9] Elzawy, M., Mosa, S., Smarandache Curves in the Galilean 4−Space G_4, Journal of the Egyptian Math. Soc., 25, 53–56, 2017.

[10] Kahraman, T., Ugurlu, H.H., Dual Smarandashe Curves and Smarandache Ruled Surfaces, Mathematical Sciences and Applications E-Notes, 2(1), 83-98,2014.

[11] Kahraman, T., Ugurlu, H.H., Smarandache Curves of a Spacelike Curve Lying on Unit Dual Lorentzian Sphere $\widetilde{S}_1^2$, CBU J. of Sci., 11(2), 93-105, 2015.

[12] Korpinar, T., Turhan, E., Smarandache $\pi_1\pi_2$ Curves of Biharmonic New Type Constant π_2−Slope Curves According to type−2 Bishop Frame in the Sol Space GDL^3, Acta Universitatis Apulensis, No.35, 245-249, 2013.

[13] Korpinar, T., New type Surfaces in terms of B−Smarandache Curves in Sol^3, Acta Scientiarum Technology, 37(3), 389-393, 2015.

[14] Korpinar, T., Turhan, E., A New Characterization of Smarandache $ts\alpha$ Curves According to Sabban Frame in Heisenberg Group $Heis^3$, Int. J. Open Problems Compt. Math., 5(4), 147-155, 2012.

[15] Kulahci, M., Bektas, M., Ergüt, M., Curves of AW(k)-Type in 3-Dimensional Null Cone, Physics Letters A 371, 275-277, 2007.

[16] Kulahci, M., Almaz, F., Some Characterizations of Osculating in the Lightlike Cone, Bol. Soc. Paran. Math., 35(2), 39-48, 2017.

[17] Liu, H., Curves in the Lightlike Cone, Contribbutions to Algebra and Geometry, 45 (1), 291-303, 2004.

[18] Liu, H., Meng, Q., Representation Formulas of Curves in a Two- and Three-Dimensional Lightlike Cone, Results Math. 59, 437-451, 2011.

[19] Mak, M., Altinbas, H., Spacelike Smarandache Curves of a Timelike Curves in Anti de Sitter 3-Space, International J. Math. Combin., Vol.3, 1-16,2016.

[20] O'Neill, B., "Semi-Riemannian Geometry with Applications to Relativity", Academic Press, London, 1983.

[21] Ozturk, U. and Ozturk, E.B.K., " Smarandache Curves According to Curves on Spacelike Surface in Minkowski 3−Space $\mathbb{R}_1^3$", Journal of Discrete Mathematics, vol. 2014, article ID829581, 10 pages, 2014.

[22] Ozturk, U. and Ozturk, E.B.K., Ilarslan, K., Nesovic, E., On Smarandache Curves Lying in Lightcone in Minkowski 3−Space, Journal of Dynamical Systems and Geometric Theories, 12(1), 81-91, 2014.

[23] Savas, M., Yakut, A.T., Tamirci, T., The Smarandache Curves on H_0^2, Gazi University Journal of Science, 29(1), 69-77, 2016.

[24] Senyurt, S., Calıskan, A., Celik, U., N^*C^*-Smarandache Curves of Bertrand Curves Pair According to Frenet Frame, International J. Math. Combin., 1, 1-7,2016.

[25] Senyurt, S., Cevahir, C., Altun, Y., On Spatial Quaternionic Involute Curve a New View, Advances in Applied Clifford Algebras, DOI: 10.1007/s00006-016-0669-7, 2016.

[26] Senyurt, S., Calıskan, A., Smarandache Curves in Terms of Sabban Frame of Fixed Pole Curve, Bol. Soc. Paran. Math., 34(2), 53-62, 2016.

[27] Senyurt, S., Calıskan, A., Smarandache Curves in Terms of Sabban Frame of Spherical Indicatrix Curves, Gen. Math. Notes, 31(2), 1-15, 2015.
[28] Taskopru, K. and Tosun, M., "Smarandache Curves on S^{2}", Boletim da Sociedade Paranaense de Matematica, 32(1), 51-59, 2014.
[29] Turgut, M. and Yilmaz, S., Smarandache Curves in Minkowski Spacetime, International Journal of Mathematical Combinatorics, vol.3, 51-55, 2008.
[30] Turhan, E., Korpinar, T., On Smarandache ts Curves of Biharmonic s-Curves According to Sabban Frame in Heisenberg group $Heis^3$, Advances Modeling and Optimization, 14(2), 343-349, 2012.
[31] Yilmaz, S., Unluturk, Y., A Note on Spacelike Curves According to Type−2 Bishop Frame in Minkowski 3−Space E_1^3, International Journal of Pure and Applied Mathematics, 103(2), 321-332, 2015.
[32] Yilmaz, S., Savcı, U.Z., Smarandache Curves and Applications According to Type−2 Bishop Frame in Euclidean 3−Space, Int. J. Math. Combin. Vol.2, 1-15, 2016.

Department of Mathematics, Firat University, 23119 ELAZIĞ/TÜRKİYE
E-mail address: mihribankulahci@gmail.com

E-mail address: fb_fat_almaz@hotmail.com